

bioGenous™ Human Colonic Organoid Kit

Catalog: K2003-HC

Product Description:

bioGenous™ Human Colonic Organoid Kit is a chemically defined cell culture medium for human colonic organoids derived from adult stem cells. Self-renewal of the colonic epithelium is driven by the proliferation of stem cells and their progenitors located in the crypts. Human colonic organoids display all hallmarks of the colonic epithelium in terms of architecture, cell type composition, and self-renewal dynamics, therefore hold great promise for unprecedented studies of human colonic development and disease, human colonic organoids may also have applications in regenerative biology through ex vivo expansion of the colonic epithelium.

Product Information:

Component	Catalog#	Volume	Storage& Stability
bioGenous™ Human Colonic Organoid Basal Medium	K2003-HC-A100/A500	100 mL/500 mL	4°C, 12 months
bioGenous™ Human Colonic Organoid Supplement B(50x)	K2003-HC-B100/B500	2 mL/10 mL	-20°C, avoid repeated freeze-thaw cycles, 12 months
bioGenous™ Human Colonic Organoid Supplement C(250x)	K2003-HC-C100/C500	0.4 mL/2 mL	-20°C, avoid repeated freeze-thaw cycles, 12 months
bioGenous™ Human Colonic Organoid Supplement D(250x)	K2003-HC-D100/D500	0.4 mL/2 mL	-20°C, avoid repeated freeze-thaw cycles, 12 months

Materials & Reagents Required But Not Included:

Manufacturer	Materials	Catalog#
bioGenous™	Primary Tissue Storage Solution	K601005
bioGenous™	Epithelial Organoid Basal Medium	B213151
bioGenous™	Organoid Dissociation Solution	E238001
bioGenous™	Anti-Adherence Rinsing Kit	E238002
bioGenous™	Organoid Cryopreservation Medium (Serum Free)	E238023
bioGenous™	Organoid Culture ECM (Reduced Growth Factor)	M315066
	Fetal Bovine Serum (FBS)	-
	DPBS (1X), liquid, contains no calcium or magnesium	-

Preparation of Human Colonic Organoid Expansion Medium and Maintenance Medium

Use a sterile technique to prepare the human colonic organoid expansion medium and maintenance medium. Human colonic organoids grown in the Human Colonic Organoid Expansion Medium overwhelmingly consisted of LGR5⁺ stem cells, cycling transit amplifying (TA) cells, early enterocytes and a small number of goblet cells. Organoids grown in Human Colonic Organoid Maintenance Medium contain LGR5⁺ stem cells, TA cells, early and mature enterocytes, goblet cells, M cells and enteroendocrine cells, as well as a low number of tuft cells. The following examples are for preparing 10 mL of Expansion Medium and Maintenance Medium. If preparing other volumes, adjust accordingly.

1. Thaw Human Colonic Organoid Supplement B (50x), Human Colonic Organoid Supplement C (250x) and Human Colonic Organoid Supplement D (250x) on ice. Mix thoroughly.
NOTE: Once thawed, use immediately or aliquot and store at -20°C for not more than 10 months. After thawing the aliquots, use immediately. Do not re-freeze.
2. For Human Colonic Organoid Expansion Medium which is used specifically for primary culture and resuscitation. Add 200 µL Human Colonic Organoid Supplement B (50x), 40 µL Human Colonic Organoid Supplement C (250x) and 40 µL Human Colonic Organoid Supplement D (250x) to 9.72 mL Human Colonic Organoid Basal Medium. Mix thoroughly.
3. For Human Colonic Organoid Maintenance Medium which is used specifically for the transfer of culture. Add 200 µL Human Colonic Organoid Supplement B(50x) and 40 µL Human Colonic Organoid Supplement C(250x) to 9.76 mL Human Colonic Organoid Basal Medium. Mix thoroughly.
NOTE: If not used immediately, store the complete medium at 2-8°C for not more than 2 weeks. bioGenous™ Human Colonic Organoid Supplement B contains fungicides and antibiotics (50x).

Protocol for Establishment of Human Colonic Organoids

CAUTION Studies involving primary human tissue material must follow all relevant institutional and governmental

regulations. Informed consent must be obtained from all subjects before the collection of the primary human tissue material.

Establishment of Organoids from Primary Tissue

1. Collect primary human colon tissue pieces in ice-cold Primary Tissue Storage Solution (K601005) with conical tubes. Keep tissue samples at 4°C until the start of the isolation.
2. Assess whether the obtained tissue pieces consist purely of epithelium or if they also contain fat or muscle tissue. If so, remove non-epithelial components as much as possible using surgical scissors or scalpels and forceps under a dissection microscope. If no fat or muscle tissue is present, continue to the next step immediately.
3. Rinse the tissue with Epithelial Organoid Basal Medium (B213151) until the supernatant is clear.
4. Thaw bioGenous™ Organoid Culture ECM (M315066) on ice and keep it cold.
5. Mince the tissue into small fragments in a cell culture dish using surgical scissors or scalpels.
CRITICAL The dissected samples must be small enough to pass through the tip of a 10 mL pipette. Digest the tissue fragments with 10mL of Tissue Digestion Solution(K601008) in a 15mL conical tube at 37°C, with a variable incubation period ranging from 50 min to 1 h. Carefully monitor the digestion process by mixing the content of the tube every 5-10 min by shaking vigorously and pipetting the mixture up and down.
CRITICAL To prevent over-digestion, one should examine the cells under the microscope if the epithelium cell clusters appear during digestion.
6. Add FBS to the tissue digestion mixture to a final concentration of 2% and filter using a 100 µm cell strainer.
7. Collect and centrifuge the filtered cells at 250g for 3 min at 4 °C. In case of a visible red pellet, aspirate the supernatant, and resuspend the pellet using 2mL of Red Blood Cell Lysis Solution (E238010) to lyse the erythrocytes at room temperature for 3 min and centrifuge at 250g for 3 min at 4°C.
8. Aspirate the supernatant and resuspend the pellet in Epithelial Organoid Basal Medium, centrifuge at 250g for 3 min at 4°C, and repeat this step one more time.
9. Aspirate the supernatant and resuspend the pellet in ECM. ECM should be kept on ice to prevent it from solidifying; thus, work quickly. The amount of ECM depends on the size of the pellet. Approximately 20-100 ducts should be plated in 25 µL of ECM.
CRITICAL Do not dilute ECM too much (ECM should be >70% (ECM vol/Total vol)) to ensure the formation of solid droplets.
10. Plate the ECM containing organoids on the bottom of 24-well cell culture plates in droplets of ~30 µL each around the center of the well.
CRITICAL Once the organoids are resuspended in ECM, proceed with plating as quickly as possible, as the ECM may solidify in the tube or pipette tip. Do not let the ECM touch the wall of the wells.
11. Place the culture plate into a humidified incubator at 37 °C and 5% (vol/vol) CO₂ for 15-25 min to let the ECM solidify.
12. Prepare the required amount of bioGenous™ Human Colon Organoid Medium.
13. Once the ECM droplets are solidified (15-25 min), open the plate and carefully add 500 µL of Organoid Complete Medium to each well.
CRITICAL Do not add the medium directly on top of the ECM droplets, as this might disrupt the droplets.
14. Place the culture plate in a humidified incubator at 37 °C and 5% (vol/vol) CO₂.
15. Change the medium every 3 days by carefully aspirating the medium from the wells and replacing it with fresh, pre-warmed Human Colon Organoid Medium.
16. Closely monitor organoid formation. Ideally, human Colon organoids should be passaged for the first time between 5 and 8 days after initial plating. Examples of successful culture in primary, passage and resuscitation of human colonic organoids are shown in Figure 1.

Splitting and Passaging of Organoids

17. Pipette up and down to disrupt the ECM, and transfer the organoid suspension into a 1.5 mL conical tube.
18. Pipette the organoid suspension up and down to mix thoroughly. Use the bottom of the tube to create pressure, which will aid the removal of ECM.
19. Centrifuge organoids at 200g for 3 min at room temperature.
20. Aspirate the supernatant, and split organoids using either mechanical disruption or Organoid Dissociation Solution (E238001). For mechanical disruption, resuspend the pellet in 1 mL of Organoid Basal Medium. Use a pipette tip to pipette the organoid suspension up and down 30 times. Use the bottom of the tube to create pressure, which will aid organoid disruption. In case of Organoid Dissociation Solution-based cell dissociation, resuspend the pellet in 0.2 mL of Organoid Dissociation Solution, pipette up and down and incubate at 37 °C until organoids fall apart. Pipette up and down with a filter tip for ≥10 times every 1 min to aid in the disruption of the organoids. Monitor digestion closely to keep the incubation time in Organoid Dissociation Solution to a minimum.
CRITICAL Do not dissociate in Organoid Dissociation Solution for >3 min, as this may result in poor organoid outgrowth or even loss of the culture. As a rule of thumb, digestion is complete if a mixture of small lumps of cells (consisting of 10–50 cells) can be observed.

21. After shearing is complete, wash once by topping up with 1 mL of Organoid Basal Medium, and centrifuge at 200g for 3 min at room temperature.
22. Aspirate the supernatant and resuspend the organoid pellet in 70% (vol/vol) ECM, and plate organoids in droplets on the bottom of a culture plate as described in Steps 12. After plating is complete, transfer the plate into a humidified incubator at 37 °C and 5% (vol/vol) CO₂ for 15–25 min.
23. Pre-warm Human Colonic Organoid Maintenance Medium at 37 °C.
24. After the ECM droplets have solidified (15–25 min), carefully pipette the pre-warmed medium into the wells.
25. Place culture plates in a humidified incubator at 37 °C and 5% (vol/vol) CO₂ until the organoids are needed for further experiments.

Appendix. Examples of different generations of human colonic organoids.




Figure 1. Examples of successful culture in primary, passage and resuscitation of human colonic organoids. (A) The growth status of human colonic organoids in primary culture (P0). The organoids are irregular folds with smooth edges and with obvious budding, and the diameter is about 80-100 μm. (B) The growth status of human colonic organoids in the first passage culture (P1), passage organoids mainly presented vesicle and with obvious budding. (C) The resuscitated cultured organoids showed a steady growth trend. (scale bar: 200 μm).

Last updated on 20th July 2023