

bioGenous™ Organoid Culture ECM (Reduced Growth Factor)

Catalog: M315066

Product Description

Basement membranes are thin continuous layers of specialized extracellular matrix that form the interface on which epithelial, endothelial, or neuronal cells grow. Basement membranes also support muscle and Schwann cells of the peripheral nerves as they are found surrounding these cells. bioGenous™ Organoid Culture ECM is a soluble form of basement membrane extracted and purified from the Engelbreth-Holm-Swarm (EHS) tumor. The major constituents of the bioGenous™ Organoid Culture ECM include laminin, collagen IV, entactin/nidogen, and heparin sulfate proteoglycan.

bioGenous™ Organoid Culture ECM is a reduced growth factor hydrogel that allows the complete control of the organoid culture microenvironment to allow for more consistent cell growth and reproducible experiments. It supports the growth and maintenance of human embryonic stem cells (hESCs), tissue-derived stem cells, and for the in vivo study of both healthy and tumor organoids derived from various organs. bioGenous™ Organoid Culture ECM is compatible with all culture media and gels at temperatures above 10°C to form a reconstituted basement membrane.

Product Information

⚠ bioGenous™ Organoid Culture ECM will start to gel if kept above 10°C for an extended period. We recommend to make aliquots and store at -80°C to -20°C to avoid repeated freeze-thaw cycles.

Component	Cat#	Volume	Concentration	Storage & Stability
bioGenous™ Organoid Culture ECM (Reduced Growth Factor)	M315066	10 mL	8.5-9.0 mg/mL	-20°C, 30 months

Directions for Use

⚠ This product is for research use only.

1. Thaw the bioGenous™ Organoid Culture ECM in a 4°C refrigerator, overnight.
2. After the bioGenous™ Organoid Culture ECM has been completely thawed, gently swirl the vial to mix the contents while keeping it on ice at all times.
3. If working with small volumes, aliquot the bioGenous™ Organoid Culture ECM by gently pipetting it into pre-chilled tubes on ice, and immediately store any remaining volume at -20°C.

Note: All cultureware, including pipettes and centrifuge tubes, should be pre-chilled or kept ice-cold to prevent the bioGenous™ Organoid Culture ECM from solidifying. If solidification occurs, the matrix can be re-liquified by placing it at 4°C on ice for 24-48 hours.

4. The volume ratio of Organoid Culture ECM should be maintained at over 70% to ensure the structural stability of the ECM during the culture process.
Note: Excessive dilution of bioGenous™ Organoid Culture ECM below 50% dilution may result in an extremely thin and fragile non-gelled protein layer that cannot support continuous organoid growth.
5. Dispense the required volumes of the ECM-cell mixture in the well.
6. Incubate the plates at 37°C for 15-20 min and then add the appropriate volume of a pre-warmed growth medium for the specific organoid type.

Directions on Coating

bioGenous™ Organoid Culture ECM (Reduced Growth Factor) could be used in the preparation of thin Gel, thick gel, or thin coating. Thin Gel preparations allow cells to be cultured on top of the gel while the Thick Gel enables the growth of cells within a three-dimensional matrix. Thin Coating could also be done to support cell growth on top of a complex protein layer.

Thin Gel

1. Thaw the required amount of bioGenous™ Organoid Culture ECM.
2. Gently mix the bioGenous™ Organoid Culture ECM, taking care to avoid introducing bubbles into the gel.
3. Add 50 $\mu\text{L}/\text{cm}^2$ bioGenous™ Organoid Culture ECM to the pre-chilled culture plate.
4. Incubate the plates at 37°C for 30 min.
5. Aspirate any unbound gel from the surface of the plate and rinse once with Basal organoid medium. Ensure that the tip of the pipet does not scratch the coated surface.

Thick Gel

1. Thaw the required amount of bioGenous™ Organoid Culture ECM.
2. Gently mix the bioGenous™ Organoid Culture ECM, taking care to avoid introducing bubbles into the gel.
3. Add the recommended number of cells to the bioGenous™ Organoid Culture ECM on ice, and dispense 150-200 $\mu\text{L}/\text{cm}^2$ onto the growth surface.
4. Incubate the plates at 37°C for 30 min.
5. Add culture medium and incubate at 37°C. Cells may also be cultured on top of this gel.

Thin Coating Method

1. Thaw the required amount of bioGenous™ Organoid Culture ECM.
2. Gently mix the bioGenous™ Organoid Culture ECM, taking care to avoid introducing bubbles into the gel.
3. Using Basal organoid medium, dilute the bioGenous™ Organoid Culture ECM. It is necessary to optimize the dilution media to determine the optimal concentration for your investigation.
4. Dispense enough diluted bioGenous™ Organoid Culture ECM to cover the entire growth surface.
5. Incubate the plates at room temperature for 60 min.
6. If any unbound materials remain, aspirate and rinse gently using the Basal Organoid Medium.
Note: *If gel-coated plates are not used immediately after coating, add PBS to prevent evaporation and store at 4°C for up to 5-10 days.*

Quality Control

All components are negative for bacterial and fungal contamination. Certificate of authenticity (COAs) for all other products are available upon request.

Safety Information

Read the Safety Data Sheets (SDSs) and follow the manufacture's instruction.

Disclaimer

To the fullest extent permitted by applicable law, bioGenous BIOTECH, Inc. and/or its affiliates shall not be liable for any special, incidental, indirect, punitive, multiple, or consequential damages arising from or related to this document or your use thereof.

Contact and Support

For questions, suggestions, and technical supports, please contact us at E-mail: info@biogenous.cn.

Last updated on 30th August, 2024